

④

FILE COPY

OFFICE OF NAVAL RESEARCH

Contract N00014-84-G-0201

Task No. 0051-865

Technical Report #24

Binuclear Phthalocyanines with Aromatic Bridges

By

H. Lam, S.M. Marcuccio, P.I. Svirskaya, S. Greenberg, A.B.P. Lever\*,  
C.C. Leznoff and R.L. Cerny

in

Canadian Journal of Chemistry

York University

Department of Chemistry, 4700 Keele St., North York  
Ontario, Canada M3J 1P3

DTIC  
ELECTE  
20 MAR 1989  
S E D

Reproduction in whole, or in part, is permitted for any purpose of the United States Government

\*This document has been approved for public release and sale; its distribution is unlimited

\*This statement should also appear in Item 10 of the Document Control Data-DD form 1473. Copies of the form available from cognizant contract administrator

AD-A205 868

89 3 20 0-18

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY Unclassified		3. DISTRIBUTION / AVAILABILITY OF REPORT As it appears on the report	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Report # 24		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION A.B.P. Lever, York University Chemistry Department	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6c. ADDRESS (City, State, and ZIP Code) 4700 Keele St., North York, Ontario M3J 1P3 Canada		7b. ADDRESS (City, State, and ZIP Code) Chemistry Division 800 N. Quincy Street Arlington, VA 22217 U.S.A.	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-84-G-0201	
8c. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Binuclear Phthalocyanines with Aromatic Bridges			
12. PERSONAL AUTHOR(S) H. Lam, S.M. Marcuccio, P.I. Svirskaya, S. Greenberg, A.B.P. Lever*, C.C. Leznoff and R.L. Cerny			
13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM Aug. 88 to Aug. 89	14. DATE OF REPORT (Year, Month, Day) March 1, 1989	15. PAGE COUNT 42
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		Phthalocyanines; Binuclear . (Mg)	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Three different binuclear phthalocyanines were prepared and characterized. Two of the binuclear phthalocyanines are cofacially linked by rigid naphthalene and anthracene spacers, while the third consists of a direct linkage between two phthalocyanine rings. The dicobalt, dicopper and dizinc derivatives were prepared by inserting the metal into the metal free species. The synthesis, physical and spectroscopic properties of the precursor bisphthalonitriles are also included.			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> OTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified/unlimited	
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Robert K. Grasselli		22b. TELEPHONE (Include Area Code)	22c. OFFICE SYMBOL

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Martin Fleischmann  
Department of Chemistry  
University of Southampton  
Southampton SO9 5H UNITED KINGDOM

Dr. John Wilkes  
Department of the Air Force  
United States Air Force Academy  
Colorado Springs, Colorado 80840-6528

Dr. R. A. Osteryoung  
Department of Chemistry  
State University of New York  
Buffalo, New York 14214

Dr. Janet Osteryoung  
Department of Chemistry  
State University of New York  
Buffalo, New York 14214

Dr. A. J. Bard  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. Steven Greenbaum  
Department of Physics and Astronomy  
Hunter College  
695 Park Avenue  
New York, New York 10021

Dr. Donald Sandstrom  
Boeing Aerospace Co.  
P.O. Box 3999  
Seattle, Washington 98124

Mr. James R. Moden  
Naval Underwater Systems Center  
Code 3632  
Newport, Rhode Island 02840

Dr. D. Rolison  
Naval Research Laboratory  
Code 6171  
Washington, D.C. 20375-5000

Dr. D. F. Shriver  
Department of Chemistry  
Northwestern University  
Evanston, Illinois 60201

Dr. Alan Bewick  
Department of Chemistry  
The University of Southampton  
Southampton, SO9 5NH UNITED KINGDOM

Dr. Edward Fletcher  
Department of Mechanical Engineering  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Bruce Dunn  
Department of Engineering &  
Applied Science  
University of California  
Los Angeles, California 90024

Dr. Elton Cairns  
Energy & Environment Division  
Lawrence Berkeley Laboratory  
University of California  
Berkeley, California 94720

Dr. Richard Pollard  
Department of Chemical Engineering  
University of Houston  
Houston, Texas 77004

Dr. M. Philpott  
IBM Research Division  
Mail Stop K 33/801  
San Jose, California 95130-6099

Dr. Martha Greenblatt  
Department of Chemistry, P.O. Box 939  
Rutgers University  
Piscataway, New Jersey 08855-0939

Dr. Anthony Sammells  
Eltron Research Inc.  
4260 Westbrook Drive, Suite 111  
Aurora, Illinois 60505

Dr. C. A. Angell  
Department of Chemistry  
Purdue University  
West Lafayette, Indiana 47907

Dr. Thomas Davis  
Polymers Division  
National Bureau of Standards  
Gaithersburg, Maryland 20899

ABSTRACTS DISTRIBUTION LIST 11/9/62

Dr. Henry S. White  
Department of Chemical Engineering  
and Materials Science  
151 Amundson Hall  
421 Washington Avenue, S.E.  
Minneapolis, Minnesota 55455

Dr. Daniel A. Buttry  
Department of Chemistry  
University of Wyoming  
Laramie, Wyoming 82071

Dr. W. R. Fawcett  
Department of Chemistry  
University of California  
Davis, California 95616

~~Dr. Peter M. Blonsky  
Eveready Battery Company, Inc.  
25225 Detroit Road, P.O. Box 45035  
Westlake, Ohio 44145~~

BINUCLEAR PHTHALOCYANINES WITH AROMATIC BRIDGES

H. Lam, S.M. Marcuccio, P. I. Svirskaya, S. Greenberg, A.B.P. Lever  
and C.C. Leznoff<sup>1</sup>

Department of Chemistry, York University, 4700 Keele Street  
North York, Ontario Canada M3J 1P3

and

Ronald L. Cerny

Midwest Center for Mass Spectrometry, University of Nebraska-  
Lincoln, Lincoln, NE 68558, U.S.A.

---

1 Author to whom correspondence may be addressed.

## ABSTRACT

Three different binuclear phthalocyanines were prepared and characterized. Two of the binuclear phthalocyanines are cofacially linked by rigid naphthalene and anthracene spacers, while the third consists of a direct linkage between two phthalocyanine rings. The dicobalt, dicopper and dizinc derivatives were prepared by inserting the metal into the metal free species. The synthesis, physical and spectroscopic properties of the precursor bisphthalonitriles are also included.

The use of porphyrins and phthalocyanines in electrocatalysis have been the subject of intensive research (1).

Four electron reductions of oxygen to water using cofacial dicobalt porphyrins have been reported by different groups of researchers (2,3). The cofacial structures have been achieved either by using a cyclophane type of framework (2) or by covalently linking two porphyrins on rigid spacers, via the 1,8-positions of anthracene and biphenylene (3). Some of these binuclear porphyrins are sensitive to light, oxygen, or both (2). To overcome this problem, we have synthesized metal free binuclear phthalocyanines covalently linked by one (4), two (5), four (5), five (6), and "-1" (7) atom bridges and some of their metallated derivatives. In general these phthalocyanines are very stable towards light and oxygen. The dicobalt derivatives of these binuclear phthalocyanines catalyze the two electron reduction of oxygen to hydrogen peroxide but the four electron process was not observed (8). The lack of a permanent cofacial interaction between the two phthalocyanine rings may be one of the reasons for not achieving the goal of a four electron reduction. We wished to prepare phthalocyanine derivatives analogous to the cofacial porphyrins which catalyze the four electron reduction of oxygen to water. Methods were developed to link phthalonitrile units to rigid spacer molecules to form the bridging bisnitriles, which can be converted to binuclear phthalocyanines.

We report herein the synthesis of three examples of binuclear phthalocyanines and their dicobalt, dicopper, and dizinc derivatives. In the direct linked binuclear phthalocyanines, the two phthalocyanine rings are joined as two phenyl groups in biphenyl derivatives, while in the cofacial systems the two phthalocyanine rings are linked via the 1,8-positions of



naphthalene (9) and anthracene. With these new binuclear phthalocyanines, we have now synthesized a series of binuclear phthalocyanines in which the covalent bridging linkages range from 5 to "-1" atoms.

#### Synthesis of Directly Linked Binuclear Phthalocyanines

Using a method previously described (10), elemental nickel was generated by reduction of nickel iodide with lithium in dry glyme in the presence of naphthalene. The activated nickel powder catalyzed the self coupling of 4-iodophthalonitrile (1) to give 3,3',4,4'-tetracyanobiphenyl (2) in 78% yield. Compound 2 was converted into its 1,3-diiminoisoindoline (3) (5,6). Condensation of 3 with 5-neopentoxo-1,3-diiminoisoindoline (4) (5,6) was carried out in 2-N,N-dimethylaminoethanol for 36 h. The resulting dark blue solution was purified by flash chromatography (11) to give the mononuclear 2,9,16,23-tetraneopentoxophthalocyanine (5) and the binuclear 2,2-bi-(9,16,23-trineopentoxophthalocyanine) (6a) in 55% and 33% yield respectively. We can refer to 6a as the zero atom bridged binuclear phthalocyanine. The binuclear phthalocyanine 6a was converted to its dicobalt 6b and dicopper 6c derivatives by refluxing 6a in toluene/2-methoxyethanol with  $\text{CoCl}_2$  and  $\text{Cu}(\text{OAc})_2$  respectively (5,6). Compound 2 and 6a-c have been fully characterized by spectroscopic data and elemental analysis (see experimental). Some spectroscopic data of 6a but not its synthesis or characterization have been previously mentioned (12).

(Scheme 1)

Synthesis of Binuclear Phthalocyanines Covalently Bridged by  
Naphthalene

Treatment of **1** and 1,8-diiodonaphthalene (**7**) with elemental nickel (**10**) at room temperature for 3.5 h gave 1,8-bis(3,4-dicyanophenyl)naphthalene (**8**), 1-iodo-8-(3,4-dicyanophenyl)naphthalene (**9**), and 3,3',4,4' tetra-cyanobiphenyl (**2**) in 15.5%, 2.5% and 42.5% yield respectively. Compound **8** was converted to its 1,3-diiminoisoindoline (**10**) as previously described (5,6). Condensation of **10** and **4** in 2-N,N-dimethylaminoethanol for 44 h gave a dark green solution. Purification by flash chromatography (11) followed by vacuum liquid chromatography (13) gave the mononuclear **5** and the binuclear **11a** in 45% and 8.7% yield respectively. The binuclear **11a** was converted to its dicobalt **11b**, dicopper **11c** and dizinc **11d** derivatives by refluxing **11a** in toluene/2-methoxyethanol with  $\text{CoCl}_2$ ,  $\text{Cu}(\text{OAc})_2$  and  $\text{Zn}(\text{OAc})_2$  respectively. Although **11a-d** and **8** have been reported in a communication (9), the full details are reported here (see experimental). The binuclear phthalocyanines **11a-d** exhibit parent ion clusters in their fast atom bombardment (FAB) mass spectra (14). More importantly, no evidence of partially or half metallated species of **11b-d** were observed. The ultra-violet-visible (UV) spectrum of **11a** gave a single broad (symmetric shape) absorption at the Q band region which has not been observed before in other metal free binuclear phthalocyanines (7,12). A more detailed  $^1\text{H}$  nmr analysis of **8** than previously reported (9) indicates that the compound exists as a mixture of two rotamers at room temperature in a 1:1 ratio. The existence of rotamers suggests that there is a rotation barrier sufficiently higher than that of the unsubstituted 1,8-biphenylnaphthalene (15). The  $^{13}\text{C}$  nmr of **8** also indicates the presence of rotamers at room temperature.

These rotamers did not separate by chromatographic methods. Although the two rotamers of **8** could be observed on the nmr time scale, their ultimate use in the formation of phthalocyanines at high temperatures, resulting in their subsequent equilibration again, removed any incentive to resort to heroic efforts to isolate the rotamers of **8**. After condensation, the binuclear phthalocyanine **11a** should also exist as two isomers due to restriction in the rotation of the phthalocyanine rings. The two possible isomers of **11a** were isolated as one fraction by chromatography. Construction of molecular models of **11a** has shown substantial overlap of the two phthalocyanine rings in each isomer, and this may be the reason that these almost identical isomers are not separable by column chromatography. Separation of these isomers of **11a** using high performance liquid chromatography (HPLC) on a silica gel column was also not successful.

(Scheme 2)

#### Synthesis of Binuclear Phthalocyanines Covalently Bridged by Anthracene

The coupling reaction of aromatic halides using elemental nickel works well only with reactive halides such as iodides. The 1,8-diiodoanthracene is unknown. Even though 1,8-dichloroanthracene is readily available (16), it is not active enough to undergo the desired cross coupling reaction.

Less electropositive arylzinc derivatives can tolerate various electrophilic functional groups such as nitriles and esters (17). These organometallic reagents readily undergo cross coupling reaction with aryl halides (18), making arylzinc derivatives the reagents of choice in our synthesis. In general, direct attack of aryl halide by zinc metal is difficult and the

organozinc reagents are usually obtained by a two step process from organolithium or Grignard reagents and the zinc halides (19). Instead of using an alkyllithium/aryl halide reaction to generate the organolithium intermediate, the organolithium can be generated conveniently by reacting the aryl halide directly with lithium metal in tetrahydrofuran (THF) using ultrasonic irradiation without any noticeable side reaction (20).

Treatment of 1,8-dichloroanthracene (12) with lithium and zinc bromide in dry THF at 0°C with ultrasonic activation for 2 h generated the 1,8-organozinc intermediate (13) (21). A cross coupling reaction between 1 and 13 catalyzed by tetrakis(triphenylphosphine)palladium (17,18) gave 1,8-bis-(3,4-dicyanophenyl)anthracene (14) in 9.9% yield. Compound 14 was converted to its 1,3-diiminoisoindoline (15) and condensed with 4 in 2-N,N-dimethyl-aminoethanol to give a dark green mixture. Upon dilution with water the mixture gave a blue colored residue. The mixture was purified by flash chromatography (11) to give a mixture of mononuclear phthalocyanine 5 and binuclear phthalocyanine 16a. A trace of the mononuclear phthalocyanine 5 was removed by gel permeation chromatography (22) to give the binuclear 16a in 12% yield. Metal free binuclear phthalocyanine 16a was converted to its dicobalt 16b and dizinc 16c derivatives by standard procedures.

Bisnitrile 14 unlike 8 does not exist as a mixture of rotamers as shown by its simpler  $^1\text{H}$  and  $^{13}\text{C}$  nmr spectra because of the low rotation barrier between the two phenyl rings in the 1,8-anthracene system (23). The UV spectrum of binuclear phthalocyanine 16a exhibits broad symmetric absorption similar to 11a. Parent ions were exhibited in the (FAB) mass spectra of 16a-c and no evidence for partially metallated species for 16b-c was observed.

Attempts to apply a similar organozinc procedure to 1,8-diiodonaphthalene did not give the desired product **8** (17,18,19). The products isolated were found to be 1-(3,4-dicyanophenyl)naphthalene (**18**) and **2**. Due to the close proximity of the 1,8-positions on naphthalene, the formation of a monozincnaphthalene intermediate **17** may be favourable over the naphthalene-1,8-dizinc species. Reaction of **1** and **17** with zinc dissociated from naphthalene after reaction may explain the formation of **18**. The elemental nickel used in the cross coupling reaction is not a true catalyst since it is not regenerated after the reaction (10). Large amounts of nickel iodide (more than 1:1 ratio) have been used for satisfactory results. Thin layer chromatography of the reaction mixture has shown that the cross coupling between the organozinc derivatives and 4-iodophthalonitrile (**1**) is faster than the homo-coupling of **1** itself. In this regard, the cross coupling reaction of organometallics and aryl halides catalyzed by transition metal complexes is superior to the mixed coupling between different aryl halides catalyzed by elemental nickel. The cross coupling product is expected to be the major product in the case of the organometallic/aryl halide reaction, whereas the homo-coupling product is expected to be the major product in the elemental nickel catalyzed aryl halides coupling. The low yield of 1,8-bis(3,4-dicyanophenyl)naphthalene (**8**) is predictable because the 1,8-diiodonaphthalene has to react twice with 4-iodophthalonitrile to form the product, and have to compete against the homo-coupling reaction. The poor yield of 1,8-bis(3,4-dicyanophenyl)anthracene which is obtained via a different coupling may be due to incomplete formation of the 1,8-dilithiated intermediate. Half of the 1,8-dichloroanthracene was recovered. Also, the possible formation of polymeric organozinc in which the zinc atom acts as a

bridge to link up adjacent anthracene molecules may decrease the yield.

(Scheme 3)

(Scheme 4)

#### Physical and Spectroscopic Properties of the Phthalonitriles

Phthalonitriles **2**, **8**, **14** and **18** all have melting points higher than their unsubstituted parent compounds (16,24) and exhibit characteristic CN absorption in the infrared region at  $2240\text{ cm}^{-1}$ . Phthalonitriles **8** and **9** have the same molecular weight of 380. High resolution mass spectra of **8** and **9** were obtained to check their elemental composition. Together with elemental analysis data, phthalonitrile **8** was clearly identified as the symmetrical 1,8-disubstituted product.

The proton nmr spectra of the bisphthalonitriles all showed complicated patterns in the aromatic region. Chemical shifts assignments were achieved for **2**, **8**, **9**, **14** and **18** with additional coupling information obtained from correlation nmr and homo-decoupling experiments (Table 1). Carbon/proton correlation experiments have been done on the more soluble phthalonitriles (Table 2). The  $^1\text{H}$  nmr and  $^{13}\text{C}$  nmr of **2** are relatively simple. Selective irradiation of  $\text{H}_2'$ ,  $\text{H}_5'$  and  $\text{H}_6'$  in **2**, which resulted in changes in signal intensity in the  $^{13}\text{C}$  spectra due to the n.o.e. effect helps to identify the chemical shifts of the carbons (26,27).

For the 1,8-disubstituted naphthalene systems, the two parallel phenyl rings are at an angle of about  $70^\circ$  to the plane of the naphthalene ring in order to relieve some of the strain in the molecule (25). The distance

between the 1,8-positions on the naphthalene ring is about 2.45 Å (24) which is too small for free rotation of the 3,4-dicyanosubstituted phenyl rings at room temperature. The  $^1\text{H}$  nmr of **8** at room temperature showed evidence that there is a sufficiently high rotation barrier between the two phenyl rings and **8** exists as two rotamers. The rotamer with two sets of cyano groups on the same side of the naphthalene plane is referred to as **8a** and the other rotamer with cyano groups on the opposite side is referred to as **8b**.

Proton/proton correlation experiments clearly established the coupling relationship of the protons in the rotamers. The correlation spectrum indicated the protons on the phenyl rings of **8a** and **8b** as separate systems (Figure 1). The chemical shifts for protons on the naphthalene ring do not seem to be affected by the different orientations of the phenyl rings. A model based on the different resultant dipoles pointed in different directions for **8a** and **8b** can help to assign the chemical shifts for the protons on the phenyl rings (Figure 2). The orientation of the resultant dipoles may influence the localization of electron density over the ring and hence the deshielding effect of the  $\pi$  cloud. For **8a**, the resultant influence located between  $\text{C}_3'$  and  $\text{C}_4'$ , which may have roughly equal effects on  $\text{H}_2'$  and  $\text{H}_5'$  and less effect on  $\text{H}_6'$ . The expected chemical shifts are therefore  $\text{H}_2' = \text{H}_5' > \text{H}_6'$ . For **8b**, the resulting influence concentrated around  $\text{C}_4'$ , affecting  $\text{H}_5'$ ,  $\text{H}_2'$  and  $\text{H}_6'$  in decreasing order. The expected chemical shifts are therefore  $\text{H}_5' > \text{H}_2' > \text{H}_6'$ . The strongest influence will be experienced by  $\text{H}_5'$  in **8b** due to the close distance between  $\text{H}_5'$  and the net dipole as shown in Figure 2. With this model, the sequence of the predicted chemical shifts of the phenyl protons are found to be matched with the observed sequence. The nitrile groups cause the phenyl protons to shift downfield which offsets

the upfield shifting effect observed in the 1,8-diphenylnaphthalene, in which the ten phenyl protons resonate at 6.85 ppm in carbon tetrachloride (24). A carbon nmr spectrum of **8** at room temperature showed 18 peaks. If only one rotamer is present, 14 peaks are expected. At 100°C in DMSO- $d_6$  the spectrum consisted of 14 peaks as expected (see experimental).

The proton spectrum of **9** is much simpler than **18** even though these compounds have similar structures. The chemical shifts of  $H_{2'}$  and  $H_{5'}$  of **9** are more downfield shifted than  $H_{6'}$ . The characteristic C-I carbon appeared at 92.07 ppm. The distance between the 1,8-positions on anthracene is about 4.8 Å apart (16). The energy barrier of rotation of the phenyl rings at the 1,8-positions is quite small (23). In fact,  $^1H$  nmr and  $^{13}C$  nmr of **14** did not indicate the presence of rotamers at room temperature. Again,  $H_{2'}$  and  $H_{5'}$  are further downfield than  $H_{6'}$  and are consistent with the chemical shifts in the naphthalene systems.

(Figure 1)

(Figure 2)

(Table 1)

(Table 2)

#### Spectroscopic Properties of the Phthalocyanines

The infrared spectra of all metal free phthalocyanines exhibited characteristic NH absorption at 3300 and 1020  $cm^{-1}$  which disappeared upon formation of their metallated derivatives. All phthalocyanines exhibited parent ions or ion clusters in their FAB mass spectra (14).



The double Q band seen in the mononuclear metal free phthalocyanines as a consequence of  $D_2h$  symmetry is further split due to coupling in the binuclear phthalocyanines (12). Coupling in general can be expected to occur either through space in the close cofacial conformation, or via conjugation through the unsaturated bridge or both. For binuclear phthalocyanines with low symmetry, transitions to higher and lower energy combinations are allowed, resulting in both blue and red shifts with respect to the mononuclear species (12). In fact, the zero linked binuclear phthalocyanine 6a and the previously reported binuclear phthalocyanines (4,5,6) except the negative linked binuclear phthalocyanine (7), show such splitting patterns. However, the electronic spectra of metal free binuclear phthalocyanines 11a and 16a which closely resembled one another, exhibited a single broad Q band absorption at 648 nm but with some evidence of splitting in the spectrum of 16a (Figure 3).

Indeed the spectra of 11a and 16a looked similar to the deprotonated spectrum of the five atom bridged binuclear phthalocyanine (12) which is able to assume a close cofacial  $D_4h$  configuration. The similarity suggests that the two phthalocyanine rings in 11a and 16a are very close together and the four internal hydrogens are equivalent which raises the symmetry of the molecules effectively to  $D_4h$ . For phthalocyanine with  $D_4h$  symmetry, the  $\pi-\pi^*$  transition is not split. There are large blue shifts in the Q band absorption in the metal free cofacial  $D_4h$  binuclear phthalocyanines 11a and 16a (Table 3) relative to mononuclear metal free phthalocyanines indicative of the extensive electronic coupling between the cofacial rings (28,29).

In general, all the metallated cofacial binuclear phthalocyanines showed two Q band absorptions, the relative intensity varying with

concentration. The absorption centered around 680 nm corresponds to the expected transition for uncoupled metallated phthalocyanines, whereas the 640 nm transition resulted from aggregated species (30). For cofacial systems, aggregation may be both intermolecular and intramolecular. Intermolecular aggregation is expected to be concentration dependent, while intramolecular aggregation should be concentration independent. When the concentration is sufficiently reduced, the spectrum becomes concentration independent and the remaining absorption near 640 nm arise from intramolecular coupling (aggregation) due to the cofacial arrangement (Figure 3).

(Figure 3)

(Table 3)

The proton nmr spectra of the binuclear phthalocyanines exhibited absorption peaks typical of the neopentoxy group and very weak aromatic signals. The NH absorption of **6a** occurred around -3 ppm. The NH absorption of the binuclear phthalocyanines **11a** and **16a** occurred at higher field. Binuclear phthalocyanine **11a** gives two sets of broad NH signals at -5 and -6 ppm. The two sets of signals may correspond to the two configurational isomers. The isomer which is exactly cofacial is related to the bisnitrile **8a**. The other isomer is related to bisnitrile **8b** in which the two rings are slightly away from each other but still have some degree of overlap in a cofacial manner. The broadening of all of the peaks is due to the positional isomers arising from the neopentyl substituents. The NH protons for the anthracene linked binuclear phthalocyanine **16a** were observed at -6.5 ppm and are much sharper. The sharpening of the signal may be due to increased

separation between the rings so that the neopentyl groups contribute less influence on the overall configuration and hence less variation in the orientation of the two phthalocyanine rings. The upfield shift of the internal NH signal is very characteristic of a cofacial configuration. It is a result of the ring current effect of the two phthalocyanine rings being held closely in a cofacial arrangement. Similar upfield shifts for NH protons in cofacial porphyrins are well documented (2,28). The NH signal disappeared in the  $^1\text{H}$  nmr spectra of the zinc derivatives of 11a and 16a.

#### EXPERIMENTAL

Matheson high purity argon was used to maintain inert atmosphere conditions. Infrared (ir) spectra were recorded on a Pye Unicam SP1000 infrared spectrophotometer using KBr discs. Nuclear magnetic resonance (nmr) spectra for protons and carbons were recorded on a Bruker AM300 nmr spectrometer. The position of signals are reported in  $\delta$  units. (The splitting of the signal are described as singlets (s), doublets (d), triplets (t), quartets (q), doublets of doublets (dd) or multiplets (m)). The  $^1\text{H}$  nmr spectra of  $10^{-4}$  M solutions of the phthalocyanines were obtained by averaging 500 - 3000 scans over the absorption range. The ultraviolet-visible spectra (UV) were recorded on a Hewlett Packard HP8451A Diode Array spectrophotometer. Mass spectra (ms) were recorded at 70eV on a VG Micromass 16F mass spectrometer in the EI mode. The FAB spectra were obtained with a Kratos MS-50 triple analyzer mass spectrometer equipped with a FAB ion source of standard Kratos design and Ion Tech atom gun. The sample was dissolved in chloroform and a microliter of the resulting solution added to a microliter of *o*-nitrophenyl octyl ether on the probe tip. The spectra of

the molecular ions of the binuclear phthalocyanines were obtained by signal averaging up to 256 scans over the appropriate mass range. The number in parenthesis after the indicated ion shows the percentage of the base peak represented by that ion. Melting points (mp) were determined using a Kofler hot stage melting point apparatus and are uncorrected. Flash chromatography was performed using silica gel of particle size 20-45 microns. All reactions, except the one which required sonication, were stirred with magnetic stirrer. Ultrasound activation was carried out using a Branson 1200 sonicator. All solvents were freshly distilled before use. Microanalyses were performed by Guelph Chemical Laboratories Ltd., Guelph, Ontario. Thin layer chromatography (tlc) was performed using silica gel G as the absorbant.

#### 3,3',4,4'-Tetracyanobiphenyl (2).

Using a method previously described (10), nickel iodide (dried for 4 h at 60-70°C/0.5 Torr) (4.53 g, 14.5 mmol), lithium (0.248g, 35.7 mmol) and naphthalene (0.186 g, 1.45 mmol) in 30 mL of fresh distilled dry glyme was stirred at room temperature for 12 h. To the nickel powder, precipitated as a bulky black slurry was added 1.0 g of (3.9 mmol) of 4-iodophthalonitrile (1) dissolved in 5 mL of dry glyme. The reaction mixture became warm (30-35°C) and after 2 h the reaction was completed (tlc benzene/acetonitrile (9:1)). The reaction mixture was poured into 100 mL of ice-water, the solid was filtered and washed twice with ice-water. The resulting product was washed very slowly with benzene to remove most of the naphthalene. It was then extracted with ethyl acetate, dichloromethane and acetonitrile (three times 30-40 mL portions for each solvent). The ethyl acetate, dichloromethane and acetonitrile solutions were combined and dried over MgSO<sub>4</sub>. The

solvent was evaporated to give 0.5 g of crude product. The crude product was chromatographed by flash chromatography (11) using benzene, benzene/-dichloromethane (1:1) and acetonitrile/dichloromethane (5:100) as eluants to give in 78% yield 0.39 g of **2** as white crystals, mp 291-293°C;

ir ( $\text{cm}^{-1}$ ): 2240 (CN), 1600, 1480, 1380, 1210, 910, 860, 840; uv ( $\text{CH}_2\text{Cl}_2$ )

$\lambda_{\text{max}}$  (log  $\epsilon$ ): 242 (4.16), 270 (4.18), 290 (4.14sh) nm;  $^1\text{H}$  nmr and  $^{13}\text{C}$  nmr:

see Tables 1 and 2; ms  $m/z$ : 254 ( $\text{M}^+$ , 100), 227 ( $\text{M}^+ - \text{HCN}$ , 38), 220

( $\text{M}^+ - 2\text{HCN}$ , 20). Anal. Calcd. for  $\text{C}_{18}\text{H}_8\text{N}_4$ : C 75.60, H 2.36, N 22.04;

found: C 75.54, H 2.07, N 21.98.

#### 2,2-Bi-(9,16,23-trineopentoxypthalocyaninyl) (6a).

By a previously described method (5,6), the two crude diiminoiso-indolines **3** and **4** were obtained from 0.273 g (1.10 mmol) of **2** and 6.0 g (28 mmol) of 4-neopentoxypthalonitrile respectively. Compounds **3** and **4** were heated at 150°C (oil bath) in 20 mL of 2-N,N-dimethylaminoethanol and 10 mL of dimethylformamide (DMF) for 36 h under an argon atmosphere. After cooling to room temperature, the dark blue mixture was diluted with water, the residue filtered and washed thoroughly with water until the filtrate was colorless. Preliminary purification of the product was achieved by flash chromatography using a 5 cm diameter column in which the crude product was preadsorbed on silica and eluted with hexane (200 mL), hexane/toluene (1:1) (1.5L) and then hexane/toluene (1:2) until all the monomeric 2,9,16,23-tetraneopentoxypthalocyanine (**5**) (3.3 g, 55% yield) was eluted. Further elution with toluene/2-methoxyethanol (200:5) (1.5 L) gave, after solvent evaporation, a portion of the dimer contaminated with traces of the monomer **5**. A second portion obtained by continued elution with toluene/2-methoxy-

ethanol (200:5) (1 L) was contaminated with some green material. The latter portion was purified by evaporating the solution to dryness, the resultant solid was suspended in dry acetone, filtered and washed with acetone until the filtrate was colorless. Finally, the product was purified by flash chromatography using silica gel and toluene/2-methoxyethanol (200:5) as eluant to give 565 mg (33% yield) of the product as a very dark, blue shining solid; ir ( $\text{cm}^{-1}$ ): 3300 (NH), 1615, 1240, 1100, 1015 (NH), 750; uv: see Table 3;  $^1\text{H}$  nmr  $\delta$ : 3.50 (br,  $\text{CH}_2\text{O}$ ), 1.40-1.20 (m,  $(\text{CH}_3)_3\text{C}$ );  $^{13}\text{C}$  nmr  $\delta$ : 31.92, 27.13, 26.49, ms  $m/z$ : 1543.8 ( $\text{M}^+$ ). Anal. Calcd. for  $\text{C}_{94}\text{H}_{94}\text{N}_{16}\text{O}_6$ : C 73.13, H 6.14, N 14.15; found: C 73.50, H 6.39, N 14.60.

2,2-Bi-(9,16,23-trineopentoxypthalocyaninyl) dicobalt (II) (6b).

A mixture of 6a (57.5 mg, 0.037 mmol), anhydrous cobalt (II) chloride (50 mg, 0.38 mmol), 2-methoxyethanol (3 mL) and toluene (7 mL) was heated at  $110^\circ\text{C}$  (oil bath) for 27 h under an argon atmosphere. The product was purified by flash chromatography using a 1.5 x 8 cm column by direct application of the refluxing mixture to the column. Elution with hot toluene gave, after solvent evaporation 24 mg of 6b. Continued elution with hot toluene/THF (1:1) yielded an additional 12 mg of 6b. The combined 36 mg of dimer 6b was washed with acetone to remove very minor fluorescent impurities (detected by tlc) leaving 32.4 mg (52% yield) of pure 6b as a very dark blue solid; ir ( $\text{cm}^{-1}$ ): 1615, 1240, 1100, 750; uv: see Table 3; ms  $m/z$ : 1657.7, ( $\text{M}^+$ ). Anal. Calcd. for  $\text{C}_{94}\text{H}_{90}\text{N}_{16}\text{O}_6\text{Co}_2$ : C 68.11, H 5.47, N 13.52, Co 7.11; found: C 68.35, H 5.86, N 13.50, Co 6.40.

2,2-Bi-(9,16,23-trineopentoxypthalocyaninyl) dicopper (II) (6c).

A mixture of **6a** (100 mg, 0.065 mmol), anhydrous copper (II) acetate (100 mg, 0.550 mmol), 2-methoxyethanol (3 mL) and toluene (7mL) was heated at 110 °C for 24 h under an argon atmosphere. The solvent was removed and the mixture continuously extracted with toluene. Evaporation of toluene under reduced pressure gave **6c** (57 mg) in 53% yield as a very dark blue shining solid; ir (cm<sup>-1</sup>): 1620, 1245, 1070, 755; uv: see Table 3; ms m/z: 1667.6, (M<sup>+</sup>). Anal. Calcd. for C<sub>94</sub>H<sub>90</sub>N<sub>16</sub>O<sub>6</sub>Cu<sub>2</sub>: C 67.73, H 5.44, N 13.44, Cu 7.62; found: C 67.72, H 5.26, N 13.15, Cu 7.55.

1,8-Bis(3,4-dicyanophenyl)naphthalene (8) and 1-iodo-8-(3,4-dicyanophenyl)-naphthalene (9).

To 30 mL of freshly distilled dimethoxyethane (DME), 23.0 g (73 mmol) of anhydrous nickel iodide, 1.0 g (144 mmol) of lithium and 0.8 g (6.25 mmol) of naphthalene were added. The mixture was stirred at room temperature for 12 h. To the resulting black slurry, 4-iodophthalonitrile (**1**) (4.0 g, 15 mmol) and 1,8-diiodonaphthalene (**7**) (25) (2.0 g, 5.26 mmol) were added. The mixture was allowed to react for 3.5 h. The resulting mixture was filtered through Celite using a sintered glass funnel and the residue was washed with ethyl acetate. The combined filtrate was washed with saturated sodium thiosulfate and water. The organic fraction was collected and dried over anhydrous magnesium sulphate. Evaporation of the solvent gave a yellowish brown solid. The crude material was purified by flash chromatography using benzene as the eluting solvent to remove naphthalene and unreacted 4-iodophthalonitrile (1.30 g). Elution with EtOAc/CHCl<sub>3</sub> (1:9) gave 50 mg of 1-iodo-8-(3,4-dicyanophenyl)naphthalene (**9**) in 2.5% yield.

Recrystallization of **9** from EtOAc/hexane gave slightly brown crystals, mp 178–180°C; ir ( $\text{cm}^{-1}$ ): 3060, 2240 (CN), 1595, 1490, 830, 780; uv ( $\text{CH}_2\text{Cl}_2$ )  $\lambda_{\text{max}}$  (log  $\epsilon$ ): 244 (4.13), 268 (4.15), 310 (4.19) nm;  $^1\text{H}$  nmr and  $^{13}\text{C}$  nmr: see Tables 1 and 2; hrms,  $m/z$  calcd. for  $\text{C}_{18}\text{H}_9\text{N}_2\text{I}$  ( $\text{M}^+$ ): 379.9809; found: 379.9802. Anal. calcd. for  $\text{C}_{18}\text{H}_9\text{N}_2\text{I}$ : C 56.86, H 2.38, N 7.37; found: C 57.21, H 2.89, N 7.89.

Further elution with the same solvent system gave 310 mg of 1,8-bis-(3,4-dicyanophenyl)naphthalene (**8**) in 15.5% yield. Recrystallization of **8** from EtOAc/hexane gave white crystals, mp 284–285°C; ir ( $\text{cm}^{-1}$ ): 3060, 2240 (CN), 1600, 1490, 835, 780; uv ( $\text{CH}_2\text{Cl}_2$ )  $\lambda_{\text{max}}$  (log  $\epsilon$ ): 238 (4.34), 265 (4.23 sh), 340 (4.04) nm; hrms,  $m/z$  calcd. for  $\text{C}_{26}\text{H}_{12}\text{N}_4$  ( $\text{M}^+$ ) 380.10619; found: 380.10580.  $^1\text{H}$  nmr and  $^{13}\text{C}$  nmr: see Tables 1 and 2;  $^{13}\text{C}$  nmr ( $\text{DMSO}-d_6$ , 100°C)  $\delta$ : 147.28, 134.67, 134.39, 134.35, 134.26, 132.52, 131.11, 130.01, 127.10, 125.22, 114.67, 114.46, 113.51, 111.62. Anal. calcd. for  $\text{C}_{26}\text{H}_{12}\text{N}_4$ : C 82.09, H 3.18, N 14.73; found: C 82.00, H 3.34, N 14.80.

Further elution with 5% acetonitrile/dichloromethane gave 850 mg of **2** in 42.5% yield.

#### Preparation of the bis-1,3-diiminoisoindoline (**10**).

The bis-1,3-diiminoisoindoline (**10**) was prepared as previously described (5,6). Bisphthalonitrile **8** (380 mg, 1.00 mmol) was added to 20 mL of a 1:1 mixture of methanol/dioxane containing 15 mg of sodium. Ammonia was bubbled into the solution at 80°C. The reaction was over in 3 h. The solution was evaporated to give crude **10** which was used directly in condensation reaction without further purification.



1,8-Bis-2'-(9',16',23'-trineopentoxypthalocyaninyl)naphthalene (11a).

The two crude diiminoisoindolines **4** and **10** obtained from 6.0 g (28.00 mmol) of 4-neopentoxypthalonitrile and 380 mg (1.00 mmol) of **8** respectively, in 25 mL of 2-N,N-dimethylaminoethanol, were heated to 150°C for 44 h. The reaction mixture was allowed to cool and then poured into 200 mL of water. The resulting mixture was filtered and washed with water until the filtrate became clear. The residue was washed again with methanol and then dried overnight in the oven at 60°C. Flash chromatography of the crude material using a 5 cm diameter column and eluting with toluene gave a mixture of mononuclear **5** and binuclear **11a**. This mixture was chromatographed further using toluene/hexane (1:1) to give 2.70 g of **5** in 45% yield. Further elution using 2-methoxyethanol/toluene (2:100) gave a fraction consisting of mainly binuclear **11a**. The final purification was achieved by vacuum liquid chromatography using toluene as the eluting solvent followed by gradual increase of 2-methoxyethanol/toluene (2:100) to give 139 mg of binuclear phthalocyanine **11a** in 8.7% yield as a dark blue solid;

ir (cm<sup>-1</sup>): 3300 (NH), 1015, 1245, 1100, 1020 (NH), 750; uv: see Table 3;

<sup>1</sup>H nmr (C<sub>6</sub>D<sub>6</sub>) δ: 7.0-8.5 (br, aromatic), 3.3-3.8 (br, CH<sub>2</sub>O), 1.0-1.6 (br, (CH<sub>3</sub>)<sub>3</sub>C), -5 to -7 (br, NH); ms m/z: 1670 (M<sup>+</sup>). Anal. calcd. for

C<sub>104</sub>H<sub>100</sub>N<sub>16</sub>O<sub>8</sub>: C 74.72, H 6.03, N 13.42; found: C 74.71, H 5.76, N 13.15.

1,8-Bis-2'-(9',16',23'-trineopentoxypthalocyaninyl)naphthalene dicobalt (II) (11b).

A mixture of 33 mg (0.02 mmol) of **11a** and 60 mg (0.46 mmol) of anhydrous cobalt (II) chloride in 10 mL of 1:4 mixture of 2-methoxyethanol/toluene was heated to 120°C for 20 h under an argon atmo-

sphere. Flash chromatography of the crude reaction mixture using a 1 cm diameter column and eluted with 2-methoxyethanol/toluene (1:50) gave 30 mg of **11b** in 85% yield as a dark blue shining solid; ir ( $\text{cm}^{-1}$ ): 2940, 1610, 1240, 1090, 750; uv: see Table 3; ms  $m/z$ : 1782 ( $M^+$ ). Anal. calcd. for  $\text{C}_{104}\text{H}_{96}\text{N}_{16}\text{O}_6\text{Co}_2$ : C 70.00, H 5.43, N 12.57, Co 6.61; found: C 70.50, H 5.50, N 13.32, Co 6.52.

1,8-Bis-2'-(9',16',23'-trineopentoxypthalocyaninyl)naphthalene dicopper (II) (**11c**).

A mixture of 53 mg (0.032 mmol) of **11a** and 57 mg (0.32 mmol) of anhydrous copper (II) acetate in 10 mL of a 1:4 mixture of 2-methoxyethanol/toluene was heated to 120°C for 20 h under argon. The crude reaction mixture was chromatographed as described above to give 37.2 mg of **11c** in 70% yield as a dark blue solid; ir ( $\text{cm}^{-1}$ ): 2940, 1610, 1235, 1090, 740; uv: see Table 3; ms  $m/z$ : 1793.6 ( $M^+ + 1$ , 100). Anal. calcd. for  $\text{C}_{104}\text{H}_{96}\text{N}_{16}\text{O}_6\text{Cu}_2$ : C 69.64, H 5.39, N 12.50, Cu 7.09; found: C 70.36, H 5.20, N 12.69, Cu 7.08.

1,8-Bis-2'-(9',16',23'-trineopentoxypthalocyaninyl)naphthalene dizinc (II) (**11d**).

A mixture of 50 mg (0.030 mmol) of **11a** and 100 mg (0.54 mmol) of anhydrous zinc acetate in 10 mL of a 1:4 mixture of 2-methoxyethanol/toluene was heated to 120°C for 20 h under argon. The crude reaction mixture was chromatographed as described for **11b** to give 44 mg of **11d** in 86% yield as a dark blue solid; ir ( $\text{cm}^{-1}$ ): 2950, 1610, 1100, 740; uv: see Table 3;  $^1\text{H}$  nmr ( $\text{CDCl}_3$ )  $\delta$ : 7.5-8.5 (br, aromatic), 3.3-4.3 (br,  $\text{CH}_2\text{O}$ ), 1.0-1.6 (br,

(CH<sub>3</sub>)<sub>3</sub>C); ms  $m/z$ : 1797.6 (M<sup>+</sup> + 1,100). Anal. calcd. for C<sub>104</sub>H<sub>96</sub>N<sub>16</sub>O<sub>6</sub>Zn<sub>2</sub>: C 69.50, H 5.38, N 12.48, Zn 7.28; found: C 69.25, H 5.73, N 12.60, Zn 7.46.

1,8-Bis(3,4-dicyanophenyl)anthracene (14).

To 60 mL of freshly distilled THF in a two neck round bottom flask, 2.0 g (8.1 mmol) of 1,8-dichloroanthracene (12) and 5.49 g (24 mmol) of anhydrous zinc bromide were added. The solution was cooled to 0°C. With a strong stream of argon flowing through the flask, 0.24 g (36 mmol) of lithium wire were cut into small pieces and dropped into the solution. The mixture was sonicated with ultrasound at 0°C for 1 h under argon and allowed to warm up to room temperature. The dark solution containing the zinc organometallic intermediate 13 was transferred with a syringe to another round bottom flask containing 4.0 g (16 mmol) of 1 and 0.8 g (4 mol %) tetrakis(triphenylphosphine)palladium. The resulting solution was stirred under argon at room temperature for 3 h. Ethyl acetate and water were added to the reaction mixture, and the cloudy mixture was filtered through Celite. The filtrate separated into an aqueous and an organic layer. The organic layer was collected and dried over anhydrous MgSO<sub>4</sub>. Evaporation of the solvent gave a brownish crude material. The crude material was purified by flash chromatography using chloroform as the eluting solvent. The first fraction contained mostly unreacted starting material. Further elution gave 34.5 mg of 1,8-bis(3,4-dicyanophenyl)anthracene (14) in 10% yield. Recrystallization of 14 from acetonitrile/EtOAc gave yellow crystals, mp 345-346°C; ir (cm<sup>-1</sup>): 3060, 2230 (CN), 1590; 1490, 880, 745; uv (CH<sub>2</sub>Cl<sub>2</sub>) λ<sub>max</sub> (log ε): 284 (4.05), 268 (4.06), 386 (3.99), 405 (3.96), ms  $m/z$ : 430 (M<sup>+</sup>, 100) 330

(20);  $^1\text{H}$  nmr and  $^{13}\text{C}$  nmr: see Tables 1 and 2. Anal. calcd. for  $\text{C}_{30}\text{H}_{14}\text{N}_4$ : C 83.69, H 3.28, N 13.02; found: C 83.42, H 3.46, N 13.15.

Preparation of the bis-1,3-diiminoisoindoline (15).

Bisphthalonitrile **14** (430 mg, 1.0 mmol) was added to 50 mL of a 1:1 mixture of methanol/dioxane containing 15 mg of sodium. Ammonia was bubbled into the solution under reflux at 80°C for 2 h. The solution was evaporated to give the crude bis-1,3-diiminoisoindoline (**15**) which was used directly in the condensation reaction without further purification.

1,8-Bis-2'-(9',16',23'-trineopentoxypthalocyaninyl)anthracene (16a).

The two crude diiminoisoindolines **15** and **4** (5,6) obtained from 430 mg (1.0 mmol) of **14** and 6.0 g (28.0 mmol) of 4-neopentoxypthalonitrile respectively in 25 mL of 2-N,N-dimethylaminoethanol, were heated to 160°C for 44 h. The reaction mixture was allowed to cool and then poured into 200 mL of water. The resulting mixture was filtered and washed with water until the filtrate became clear. The residue was washed again with MeOH and then dried in the oven at 60°C overnight. Flash chromatography of the crude material using toluene as eluting solvent gave a mixture of mononuclear **5** and binuclear **16a**. Rechromatography of the mixture using toluene/hexane (1:1) did not give a good separation. The binuclear fractions were still contaminated with large amounts of **5**. The mixed binuclear fraction was rechromatographed using gel permeation column (Bio-Beads SX-1, 200-450 mesh) with THF as the eluting solvent. The front running binuclear fractions were combined and rechromatographed on a silica gel column using 2-methoxy-ethanol/toluene (2:100) as the eluting solvent to give 210 mg of 1,8-bis-2'-

(9',16',23'-trineopentoxypthalocyaninyl)anthracene (**16a**) in 12% yield as a dark blue solid; ir ( $\text{cm}^{-1}$ ): 3300(NH), 1610, 1240, 1100, 1020 (NH), 750; uv: see Table 3;  $^1\text{H}$  nmr ( $\text{C}_6\text{D}_6$ )  $\delta$ : 7.0-8.5 (br, aromatic), 3.5-3.7 (br,  $\text{OCH}_2$ ), 1.0-1.6 (br,  $(\text{CH}_3)_3\text{C}$ ), -6 to -7 (NH); ms  $m/z$ : 1719 ( $\text{M}^+$  100%); Anal. calcd. for  $\text{C}_{108}\text{H}_{102}\text{N}_{16}\text{O}_6$ : C 75.39, H 5.98, N 13.03; found: C 75.18, H 6.33, N 12.65.

1,8-Bis-2'-(9',16',23'-trineopentoxypthalocyaninyl)anthracene dicobalt (II)  
(16b).

A mixture of 52 mg (0.030 mmol) of **16a** and 60 mg (8.46 mmol) of anhydrous cobalt (II) chloride in 10 mL of a 1:4 mixture of 2-methoxyethanol/toluene was heated to 120°C for 20 h. The crude reaction mixture was flash chromatographed using 2-methoxyethanol/toluene (1:80) and gave 56 mg of **16b** in 95% yield; ir ( $\text{cm}^{-1}$ ): 2940, 1610, 1240, 1090, 750; uv: see Table 3; ms  $m/z$ : 1832 ( $\text{M}^+$ ). Anal. calcd. for  $\text{C}_{108}\text{H}_{98}\text{N}_{16}\text{O}_6\text{Co}_2$ : C 70.71, H 5.38, N 12.23, Co 6.43; found: C 70.90, H 5.88, N 11.83, Co 6.40.

1,8-Bis-2'-(9',16',23'-trineopentoxypthalocyaninyl)anthracene dizinc (II)  
(16c).

A mixture of 56 mg (0.0325 mmol) of **16a** and 100 mg (0.54 mmol) of anhydrous zinc acetate in 10 mL of a 1:4 mixture of 2-methoxyethanol/toluene was heated to 120°C for 20 h. The crude reaction mixture was flash chromatographed using 2-methoxyethanol/toluene (1:50) to give 58 mg of **16c** in 97% yield; ir ( $\text{cm}^{-1}$ ): 2950, 1610, 1240, 1090, 750; uv: see Table 3;  $^1\text{H}$  nmr ( $\text{C}_6\text{D}_6$ )  $\delta$ : 7.5-8.5 (br, aromatic), 3.3-4.3 (br,  $\text{CH}_2\text{O}$ ), 1.0-1.6 (br  $(\text{CH}_3)_3\text{C}$ ). ms  $m/z$ : 1847 ( $\text{M}^+$ ). Anal. calcd. for  $\text{C}_{108}\text{H}_{98}\text{N}_{16}\text{O}_6\text{Zn}_2$ : C 70.22, H 5.35, N

12.14, Zn 7.08; found: C 70.72, H 5.65, N 11.60, Zn 7.45.

1-(3,4 dicyanophenyl)naphthalene (18).

Following a procedure described before, 1.0 g (2.63 mmol) of 1,8-diiodonaphthalene and 1.78 g (7.9 mmol) of anhydrous zinc bromide were added to a two necked flask containing 60 mL of freshly distilled THF. The solution was cooled to 0°C. Lithium (0.09 g, 0.013 mol) was cut into small pieces and added into the flask flushed with a vigorous stream of argon. The mixture was sonicated with ultrasound at 0°C for 1 h under argon and allowed to warm to room temperature. The resulting solution was transferred with a syringe to another round bottom flask containing 1.33 g (5.26 mmol) of **1** and 0.3 g (~ 4 mol %) tetrakis(triphenylphosphine)palladium. After stirring this mixture at room temperature for 3 h, water was added followed by ethyl acetate. The cloudy mixture was filtered through Celite. The filtrate separated into aqueous and organic layers. The organic layer was collected and dried over anhydrous MgSO<sub>4</sub>. Evaporation of the solvents gave a brownish colored crude material. The crude material was purified by flash chromatography using chloroform. The first fraction contained mostly starting material. Further elution gave 200 mg of **18** in 30% yield as white crystals, mp 148-149°C; ir (cm<sup>-1</sup>): 3060, 2230 (CN), 1600, 1485, 1390, 910, 770; uv (CH<sub>2</sub>Cl<sub>2</sub>) λ<sub>max</sub> (log ε): 242 (3.67), 282 (3.42), 324 (3.41) nm; <sup>1</sup>H nmr and <sup>13</sup>C nmr: see Tables 1 and 2; ms m/z: 254 M<sup>+</sup> (100%), 233 (15%).

Anal. calcd. for C<sub>18</sub>H<sub>10</sub>N<sub>2</sub>: C 83.69, H 3.28, N 13.02; found: C 83.42, H 3.46, N 13.15.

ACKNOWLEDGEMENTS

Financial support by the Natural Sciences and Engineering Research Council of Canada and The Office of Naval Research (Washington) is gratefully acknowledged. The work was also funded by the Midwest Center for Mass Spectrometry, a National Science Foundation Regional Instrumentation Facility. We would like to thank Mr. Byron Bachelor for his assistance in the preparation of compound 2.

## REFERENCES

1. D. Lexa, P. Millard, M. Momenteau, and J. M. Saveant. *J. Am. Chem. Soc.* **106**, 6321 (1984); P. Sayer, M. Gouterman, and C.R. Connell. *Acc. Chem. Res.* **15**, 73 (1982); P. Forswey, T. Kuwana, N. Kobayashi, and T. Osa. *Adv. Chem. Ser.* **201**, 601 (1982); J. Zagal, P. Binora, and E. Yeager. *J. Electrochem. Soc.* **127**, 1506 (1980); F. Van Den Brink, E. Barendrecht, and W. Visscher. *Trav. Chim. Pays-Bas* **99**, 253 (1980); K. Kasuea, and M. Tsutsui. *Coord. Chem. Rev.* **32**, 67 (1980).
2. J.P. Collman, P. Denisevich, Y. Konai, M. Marrocco, C. Koval, and F.C. Anson. *J. Am. Chem. Soc.* **102**, 6027 (1980); J.P. Collman, F.C. Anson, S. Bencosme, A. Chong, T. Collins, P. Denisevich, E. Evitt, T. Geiger, J.A. Ibers, G. Jameson, Y. Konai, C. Koval, K. Meier, P. Oakley, R.B. Pettman, E. Schmittou, and J. Sessler. In *Organic Synthesis Today and Tomorrow*. Edited by B.M. Trost and C.R. Hutchinson. Pergamon, Oxford. 1981; R.R. Durand, C.S. Bencosme, J.P. Collman, and F.C. Anson. *J. Am. Chem. Soc.* **105**, 2710 (1983); J.P. Collman, F.C. Anson, C.E. Barnes, C.S. Bencosme, T. Geiger, E.R. Evitt, R.P. Kreh, K. Meier, and R.B. Pettman. *J. Am. Chem. Soc.* **105**, 2694 (1983).
3. C.K. Chang and I. Abdalmuhdi. *J. Org. Chem.* **48**, 5388 (1983); C.K. Chang, H.Y. Liu, and I. Abdalmuhdi. *J. Am. Chem. Soc.* **106**, 2725 (1984); C.K. Chang and I. Abdalmuhdi. *Angew. Chem.* **23**, 164 (1984); H.Y. Liu, I. Abdalmuhdi, C.K. Chang, and F.C. Anson. *J. Phys. Chem.* **89**, 665 (1985); J.P. Fillers, K.G. Ravichandran, I. Abdalmuhdi, A. Tulinsky, and C.K. Chang. *J. Am. Chem. Soc.* **108**, 417 (1986).
4. S. Greenberg, S.M. Marcuccio, C.C. Leznoff, and K.B. Tomer. *Synthesis*, 406 (1986).
5. S.M. Marcuccio, P.I. Svirskaya, S. Greenberg, A.B.P. Lever, C.C. Leznoff, and K.B. Tomer, *Can. J. Chem.* **63**, 3057 (1985).
6. C.C. Leznoff, S. Greenberg, S.M. Marcuccio, P.C. Minor, P. Seymour, A.B.P. Lever, and K.B. Tomer. *Inorg. Chem. Acta*, **89**, L35 (1984); C.C. Leznoff, S.M. Marcuccio, S. Greenberg, A.B.P. Lever, and K.B. Tomer. *Can. J. Chem.* **63**, 623 (1985).
7. C.C. Leznoff, H. Lam, S.M. Marcuccio, W.A. Nevin, P. Janda, N. Kobayashi, and A.B.P. Lever. *J. Chem. Soc. Chem. Commun.* 699 (1987).
8. M.R. Hempstead, A.B.P. Lever, and C.C. Leznoff. *Can. J. Chem.* **65**, 2677 (1987).
9. C.C. Leznoff, H. Lam, W.A. Nevin, N. Kobayashi, P. Janda, and A.B.P. Lever. *Angew Chem. Int. Ed. Engl.* **26**, 1021 (1987).
10. S. Inaba, H. Matsumoto, and R.D. Rieke. *Tet. Lett.* **23**, 4215 (1982); H. Matsumoto, S. Inaba, and R.D. Rieke. *J. Org. Chem.* **48**, 840 (1983); S. Inaba, H. Matsumoto, and R.D. Rieke. *J. Org. Chem.* **49**, 2093 (1984).



11. W.C. Still, M. Kahn, and A. Mitra. *J. Org. Chem.* **43**, 2923 (1978).
12. E.S. Dodsworth, A.B.P. Lever, P. Seymour, and C.C. Leznoff. *J. Phys. Chem.* **89**, 5698 (1985).
13. N.M. Targett, J.P. Kilcoyne, and B. Green. *J. Org. Chem.* **44**, 4962 (1979).
14. M. Barber, R.S. Bordoli, R.D. Sedgwick, and A.N. Tyler. *J. Chem. Soc. Chem. Commun.* 325 (1981).
15. H.O. House, W.J. Campbell, and M. Gall. *J. Org. Chem.* **35**, 1815 (1970).
16. H.O. House, D. Koepsell, and W. Jaeger. *J. Org. Chem.* **38**, 1167 (1973).
17. E. Negishi. *Acc. Chem. Res.* **15**, 340 (1982); E. Negishi. In *Current Trends in Organic Synthesis*. Pergamon Press, Oxford. 1982. p. 269.
18. E. Negishi, A.O. King, and N. Okukdo. *J. Org. Chem.* **42**, 1821 (1977); T.R. Bailey. *Tet. Lett.* **27**, 4407 (1986).
19. B.J. Wakefield. In *The Chemistry of Organolithium Compounds*. Pergamon Press, Oxford. 1974. p.247.
20. J.L. Luche, C. Petrier, J.P. Lansard, and A.E. Greene. *J. Org. Chem.* **48**, 3837 (1983).
21. C. Petrier, J.C. de Souza Barbosa, C. Dupuy, and J.L. Luche. *J. Org. Chem.* **50**, 5761 (1985); P. Boudjouk, R. Sooriyakumaran, and B.H. Han. *J. Org. Chem.* **51**, 2818 (1986); D. Bremner, *Chem. in Britan*, 633 (1986); T.J. Mason. *Education in Chemistry*, 102 (1987).
22. J.A. Anton, J. Kwong and P.A. Loach, *J. Heterocycl. Chem.* **13**, 717 (1976); W.A. Nevin, W. Liu, S. Greenberg, M.R. Hempstead, S.M. Marcuccio, M. Melnik, C.C. Leznoff, and A.B.P. Lever. *Inorg. Chem.* **26**, 891 (1987).
23. H.O. House, J.A. Hrabie, and D. VanDerveer. *J. Org. Chem.* **51**, 921 (1986).
24. H.O. House, R.W. Magin, and H.W. Thompson. *J. Org. Chem.* **28**, 2403 (1963); V. Balasubramanian. *Chem. Rev.* **66**, 567 (1966).
25. H.O. House, D.G. Koepsell, and W.J. Campbell. *J. Org. Chem.* **37**, 1003 (1972).
26. J.D. Memory, and N.K. Wilson. In *MNR of Aromatic Compounds*. John Wiley & Sons, New York 1982. p. 78. F. Toda, T. Oshima, Y. Ishida, Y. Takehira, K. Saito, and K. Tanaka. In *Handbook of <sup>13</sup>C NMR Spectra*. Sankyo, Japan (1981).

27. P.E. Hansen. *Org. Mag. Res.* **12**, 109 (1979); B.P. Cho and R.G. Harvey. *J. Org. Chem.* **52**, 5679 (1987).
28. D. Dolphin, J. Hiom, and J.B. Paine III. *Heterocycles*, **16**, 417 (1981).
29. C.K. Chang. *J. Chem. Soc. Chem. Commun.* 800 (1977).
30. W.A. Nevin, W. Liu, and A.B.P. Lever. *Can. J. Chem.* **65**, 855 (1987).

Table 1.  $^1\text{H}$  nmr Chemical Shifts in ppm ( $\delta$ ) of some Phthalonitriles

Compound	Proton <sup>a</sup>						
	2'	5'	6'	2	3	4	5
2 (acetone- $d_6$ )	8.09 (d, J=1.7)	8.09 (d, J=7.9)	8.13 (dd, J=7.9)				
8a (DMSO- $d_6$ )	7.82 (d, J=1.5)	7.77 (d, J=8.1)	7.54 (d, J=8.1)	7.48 (dd, J=7.2)	7.73 (t, J=7.7)	8.23 (d, J=7.4)	7.73 (t, J=7.7)
8b (DMSO- $d_6$ )	7.72 (d, J=1.5)	7.91 (d, J=8.1)	7.61 (d, J=8.1)	7.48 (dd, J=7.2)	7.73 (t, J=7.7)	8.23 (dd, J=7.4)	7.73 (t, J=7.7)
9 (DMSO- $d_6$ )	8.19 (d, J=1.6)	8.23 (d, J=8.1)	7.86 (d, J=8.1)	7.52 (d, J=7.0)	7.63 (t, J=7.6)	8.10 (d, J=8.2)	7.27 (t, J=7.6)
14 ( $\text{CD}_3\text{CN}$ ) <sup>b</sup>	8.06 (d, J=1.8)	8.04 (d, J=8.3)	7.06 (d, J=8.1)	7.51 (d, J=7.4)	7.63 (dd, J=6.9)	8.22 (d, J=8.7)	7.63 (dd, J=6.9)
18 (DMSO- $d_6$ ) <sup>c</sup>	8.29 (d, J=1.5)	8.26 (d, J=8.1)	8.03 (d, J=8.1)	7.52 (d, J=7.1)	7.63 (t, J=7.2)	8.06 (d, J=7.2)	7.54 (t, J=7.5)

<sup>a</sup> The numbering of the carbon atoms in the compounds listed follows that given for these structures in the schemes and does not follow from the name of the compounds.

<sup>b</sup> The chemical shift values for protons 9 and 10 are 8.74 (s) and 8.13 (s) respectively.

<sup>c</sup> The chemical shift value for proton 8 is 7.71 (d, J=8.4).

Table 2.  $^{13}\text{C}$  nmr Chemical Shifts in ppm ( $\delta$ ) of Some Bisphthalonitriles and Related Compounds in  $\text{DMSO}-d_6$

Compound	Carbon <sup>a</sup>																	
	1'	2'	3'	4'	5'	6'	7'	8'	1	2	3	4	4a	5	6	7	8	8a
2	141.18	132.79	114.88 <sup>b</sup>	115.58 <sup>b</sup>	134.64	132.44	115.61 <sup>c</sup>	115.66 <sup>c</sup>	-	-	-	-	-	-	-	-	-	-
7 <sup>d</sup>	-	-	-	-	-	-	-	-	96.04	145.07	128.06	132.12	136.91	132.12	128.06	145.07	96.04	132.99
8	147.59	134.98	113.56 113.81 <sup>b,c</sup>	117.60 118.81 <sup>b,c</sup>	133.17	134.75	115.26 115.13 <sup>c,e</sup>	115.54 <sup>c</sup>	135.07	131.80	125.84	130.61	134.66	130.61	125.84	131.80	135.07	127.61
9	147.00	136.14	113.20 <sup>b</sup>	113.94 <sup>b</sup>	133.46	135.98	116.03 <sup>c</sup>	116.16 <sup>c</sup>	137.22	131.46	125.41	130.79	135.26	129.98	127.55	142.54	92.07	130.35
12 <sup>f</sup>	-	-	-	-	-	-	-	-	130.71	127.83	126.24	126.68	132.21	126.68	126.24	127.83	130.71	128.62
14 <sup>g</sup>	145.02	134.83	113.51 <sup>b</sup>	115.11 <sup>b</sup>	135.29	133.72	115.73 <sup>c</sup>	115.88 <sup>c</sup>	135.86	127.82	125.72	129.55	131.29	129.55	125.72	127.82	135.86	128.40
18	145.90	135.21	113.71 <sup>b</sup>	115.40 <sup>b</sup>	134.44	135.52	116.23 <sup>c</sup>	116.39 <sup>c</sup>	136.11	127.99	125.94	129.69	133.65	128.92	127.59	126.81	124.75	130.28

<sup>a</sup> The numbering of the carbon atoms in the compounds listed follows that given for these structures in the schemes and does not follow from the names of the compounds.

<sup>b</sup> These values may be interchanged.

<sup>c</sup> These values may be interchanged.

<sup>d</sup> Data taken from Ref. 27.

<sup>e</sup> The two given values correspond to the two tautomers 8<sub>a</sub> and 8<sub>b</sub> and may be interchanged.

<sup>f</sup> The chemical shift values for carbons 9 and 10 are 128.45 and 119.00 respectively.

<sup>g</sup> The chemical shift values for carbons 9 and 10 are 128.08 and 120.82 respectively.

Table 3. Absorption Spectra of Binuclear Phthalocyanines in  
o-dichlorobenzene at room temperature <sup>a</sup>

Pc	$\lambda_{\text{max}}$ (nm) ( $\epsilon$ log)					
6a	298 (4.79)	342 (5.94)	648 (4.78)	682 (4.89)	704 (4.90)	
6b	300 (4.90)	332 (4.90)	652 (4.79)			
6c	300 (4.76)	334 (4.84)	642 (4.74)			
11a		340 (4.90)	648 (4.89)			
11b	306 (5.15)		638 (5.09)	674 (5.01)		
11c	300 (4.88)	338 (5.01)	640 (5.02)	688 (5.02)		
11d	298 (4.88)	340 (5.05)	640 (5.03)	686 (4.96)		
16a		342 (5.01)	646 (5.00)			
16b	298 (5.03)	326 (4.97)	640 (4.94)	672 (4.95)		
16c	298 (4.71)	344 (5.01)	640 (4.93)	684 (5.04)		

<sup>a</sup> The concentration of the binuclear phthalocyanines were  $\sim 1 \times 10^{-5}$  M.

Figure 1.  $^1\text{H}$  Homonuclear Correlation (COSY) of a mixture of **8a** and **8b** in DMSO- $d_6$  at 25°C.





FIGURE 1


Figure 2. 1,8-Bis(3,4-dicyanophenyl)naphthalene rotamers **8a** and **8b**.





$$\delta \text{ H}_{2'} \approx \text{H}_{5'} > \text{H}_{6'}$$

8a



$$\delta \text{ H}_{5'} > \text{H}_{2'} > \text{H}_{6'}$$

8b

FIGURE 2

Figure 3. Absorption spectra of metal free and metallated binuclear  
phthalocyanines in o-dichlorobenzene at room temperature ([PcM] -----  
~  $1 \times 10^{-5}$  M, ..... ~  $5 \times 10^{-7}$  M).







FIGURE 2




SCHEME 1



SCHEME 2



SCHEME 3



SCHEME 4